

T violation in $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decays with polarized Λ

Chuan-Hung Chen^a, C. Q. Geng^{b,c} and J. N. Ng^c

^aInstitute of Physics, Academia Sinica

Taipei, Taiwan 115, Republic of China

^bDepartment of Physics, National Tsing Hua University

Hsinchu, Taiwan, Republic of China

^cTheory Group, TRIUMF, 4004 Wesbrook Mall

Vancouver, B.C. V6T 2A3, Canada

Abstract

We study the T violating effects in the baryonic decays of $\Lambda_b \rightarrow \Lambda l^+ l^-$ ($l = e, \mu$) with polarized Λ . We show that the transverse Λ polarizations in these baryonic decays could be as large as 50% in CP violating theories beyond the standard model such as the SUSY models, which can be tested in various future hadron colliders.

Recently, time-reversal violation (TV) has been measured experimentally in the K^0 system [1], and thus complements the information on CP violation (CPV) that has been steadily accumulating for the past thirty seven years. The data is in accordance with the CPT theorem which is fundamental to local quantum field theories with Lorentz invariance and the usual spin-statistics connection. However, the origin of the violation remains unclear. In the standard model, CPV or TV arises from a unique physical phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [2]. This paradigm also predicts CPV effect in the b-quark system. To test the accuracy of this paradigm and to search for other sources of CPV one needs to look for new processes, especially the B^0 system. Indeed this is an important quest of the B-factories. In addition we deem it particularly interesting if the time reversal symmetry violation can be directly detected in the b-system rather than inferring it as a consequence of CPT invariance. In this paper, we will study the T violating effects in the baryonic decays of $\Lambda_b \rightarrow \Lambda l^+ l^-$.

It is known that for a general three-body decay of a baryon the triple spin-momentum correlations, such as $\vec{s} \cdot (\vec{p}_i \times \vec{p}_j)$, are T-odd observables, where $\vec{s}$ and $\vec{p}_{i,j}$ are the spin and momentum vectors of the final particles, respectively. There are a number of different sources that might give rise to these T-odd observables. The most important ones being the weak CPV such as the CKM phase of the SM. However, final state interactions such as QCD for non-leptonic decays or the electromagnetic (EM) interaction among the final state particles can also make contributions. These are usually less interesting and they could even hide the signals from the weak CPV. We note that the T-odd triple correlations do not need non-zero strong phases unlike some of CP violating observables, such as the rate asymmetry between a particle and its antiparticle.

In $\Lambda_b \rightarrow \Lambda l^+ l^-$ decays, due to experimental concerns only one polarization state can be measured. We can use either the polarization of the lepton (s_l) or the Λ baryon (s_Λ) to study the T-odd correlations. For the case with a polarized lepton, since the T-odd correlation such as the transverse lepton polarization is always associated with the lepton mass, we expect that the T violating effects are small for the light lepton modes [3]. Although the τ mode has less suppression due to its mass, the corresponding branching ratio which is $O(10^{-7})$ is about one order of magnitude smaller than that for the e and μ ones. The above considerations plus the fact that the efficiency of spin measurements in general are not high make lepton polarization a poor choice. In our following analysis, we will concentrate on

the search for the possibility of large T-odd term such as $\vec{s}_\Lambda \cdot (\vec{p}_{l^+} \times \vec{p}_\Lambda)$ in $\Lambda_b \rightarrow \Lambda l^+ l^-$ with $l = e$ and μ , and we can set $m_l = 0$ within the accuracy of our calculations.

We start with the effective Hamiltonian for $b \rightarrow sl^+ l^-$ by including the right-handed couplings in the hadronic sector are given by

$$\mathcal{H}(b \rightarrow sl^+ l^-) = \frac{G_F \alpha_{em}}{\sqrt{2}\pi} V_{tb} V_{ts}^* [H_{1\mu} L_V^\mu + H_{2\mu} L_A^\mu] \quad (1)$$

with

$$\begin{aligned} H_{1\mu} &= \bar{s} \gamma^\mu (C_9^L P_L + C_9^R P_R) b - \frac{2m_b}{q^2} \bar{s} i \sigma_{\mu\nu} q^\nu (C_7^L P_R + C_7^R P_L) b, \\ H_{2\mu} &= \bar{s} \gamma^\mu (C_{10}^L P_L + C_{10}^R P_R) b, \\ L_V^\mu &= \bar{l} \gamma^\mu l, \\ L_A^\mu &= \bar{l} \gamma^\mu \gamma_5 l, \end{aligned} \quad (2)$$

where C_i^L and C_i^R ($i = 7, 9, 10$) denote the effective Wilson coefficients of left- and right-handed couplings, respectively. In the standard model,

$$\begin{aligned} C_9^L &= C_9^{eff}, \quad C_9^R = 0, \\ C_{10}^L &= C_{10}, \quad C_{10}^R = 0, \\ C_7^L &= C_7^{eff}, \quad C_7^R = \frac{m_s}{m_b} C_7^{eff}, \end{aligned} \quad (3)$$

where C_9^{eff} , C_{10} , and C_7^{eff} are the standard Wilson coefficients [4].

To study the exclusive decays of $\Lambda_b \rightarrow \Lambda l^+ l^-$, one needs to know the form factors in the transition of $\Lambda_b(p_{\Lambda_b}) \rightarrow \Lambda(p_\Lambda)$, parametrized generally as follows:

$$\begin{aligned} \langle \Lambda | \bar{s} \Gamma_\mu b | \Lambda_b \rangle &= f_1^{(T)} \bar{u}_\Lambda \gamma_\mu u_{\Lambda_b} + f_2^{(T)} \bar{u}_\Lambda i \sigma_{\mu\nu} q^\nu u_{\Lambda_b} + f_3^{(T)} q_\mu \bar{u}_\Lambda u_{\Lambda_b}, \\ \langle \Lambda | \bar{s} \Gamma_\mu \gamma_5 b | \Lambda_b \rangle &= g_1^{(T)} \bar{u}_\Lambda \gamma_\mu \gamma_5 u_{\Lambda_b} + g_2^{(T)} \bar{u}_\Lambda i \sigma_{\mu\nu} q^\nu \gamma_5 u_{\Lambda_b} + g_3^{(T)} q_\mu \bar{u}_\Lambda \gamma_5 u_{\Lambda_b}, \end{aligned} \quad (4)$$

where $\Gamma_\mu = \gamma_\mu$ ($i \sigma_{\mu\nu}$), and $f_i^{(T)}$ and $g_i^{(T)}$ are the form factors of vector (tensor) and axial-vector (axial-tensor) currents, respectively. In the heavy quark effective theory (HQET) [5], the form factors in Eq. (4) can be simplified by using

$$\langle \Lambda(p_\Lambda) | \bar{s} \Gamma b | \Lambda_b(p_{\Lambda_b}) \rangle = \bar{u}_\Lambda (F_1(q^2) + \not{q} F_2(q^2)) \Gamma u_{\Lambda_b}, \quad (5)$$

where Γ denotes the Dirac matrix, $v = p_{\Lambda_b}/M_{\Lambda_b}$ is the four-velocity of Λ_b , and $q = p_{\Lambda_b} - p_{\Lambda}$ is the momentum transfer, and the relations among the form factors can be found in Ref. [6]. Explicitly, under the HEQT, we have

$$\begin{aligned} f_1 &= g_2 = f_2^T = g_2^T = F_1 + \sqrt{r}F_2, \\ \rho &\equiv M_{\Lambda_b} \left(\frac{f_2 + g_2}{f_1 + g_1} \right) = \frac{M_{\Lambda_b}}{q^2} \left(\frac{f_1^T + g_1^T}{f_1 + g_1} \right) = \frac{F_2}{F_1 + \sqrt{r}F_2}. \end{aligned} \quad (6)$$

In order to study the T violating effects using the Λ spin polarization, we write the Λ four-spin vector in terms of a unit vector, $\hat{\xi}$, along the Λ spin in its rest frame, as

$$s_0 = \frac{\vec{p}_{\Lambda} \cdot \hat{\xi}}{M_{\Lambda}}, \quad \vec{s} = \hat{\xi} + \frac{s_0}{E_{\Lambda} + M_{\Lambda}} \vec{p}_{\Lambda}, \quad (7)$$

and choose the unit vectors along the longitudinal, normal, transverse components of the Λ polarization, to be

$$\begin{aligned} \hat{e}_L &= \frac{\vec{p}_{\Lambda}}{|\vec{p}_{\Lambda}|}, \\ \hat{e}_N &= \frac{\vec{p}_{\Lambda} \times (\vec{p}_{l^-} \times \vec{p}_{\Lambda})}{|\vec{p}_{\Lambda} \times (\vec{p}_{l^-} \times \vec{p}_{\Lambda})|}, \\ \hat{e}_T &= \frac{\vec{p}_{l^-} \times \vec{p}_{\Lambda}}{|\vec{p}_{l^-} \times \vec{p}_{\Lambda}|}, \end{aligned} \quad (8)$$

respectively. Hence, the differential decay rates with polarized Λ is given by

$$d\Gamma = \frac{1}{2} d\Gamma^0 \left[1 + \vec{P} \cdot \hat{\xi} \right], \quad (9)$$

$$d\Gamma^0(t) = \frac{G_F^2 \alpha_{em}^2 \lambda_t^2}{96\pi^5} M_{\Lambda_b}^5 \sqrt{t^2 - r} f_1^2 R_{\Lambda_b}(t) dt, \quad (10)$$

where $\vec{P}$ is the Λ polarization vector, defined by

$$\vec{P} = P_L \hat{e}_L + P_N \hat{e}_N + P_T \hat{e}_T, \quad (11)$$

and

$$\begin{aligned} R_{\Lambda_b}(t) &= \left[(3(1+r)t - 2r - 4t^2) + s\rho^2 (3(1+r)t - 4r - 2t^2) + 6\rho\sqrt{r}s(1-t) \right] \\ &\times \left(|C_9^R|^2 + |C_9^L|^2 + |C_{10}^R|^2 + |C_{10}^L|^2 \right) \\ &+ \frac{4\hat{m}_b}{s} (3(1+r)t - 4r - 2t^2) \left(|C_7^R|^2 + |C_7^L|^2 \right) \\ &- 6\sqrt{r}s [1 + 2(t-r)\rho + s\rho^2] (\text{Re}C_9^R C_9^{L*} + \text{Re}C_{10}^R C_{10}^{L*}) \\ &+ 12\hat{m}_b [2\rho(r+t^2 - (1+r)t) - \sqrt{r}(1-t)(1+s\rho^2)] (\text{Re}C_9^R C_7^{L*} + \text{Re}C_9^L C_7^{R*}) \\ &+ 12\hat{m}_b [2\rho\sqrt{r}s + (t-r)(1+s\rho^2)] (\text{Re}C_9^R C_7^{R*} + \text{Re}C_9^L C_7^{L*}), \end{aligned} \quad (12)$$

with $\lambda_t = V_{tb}V_{ts}^*$, $t = E_\Lambda/M_{\Lambda_b}$, $r = M_\Lambda^2/M_{\Lambda_b}^2$, $\hat{m}_b = m_b/M_{\Lambda_b}$, and $s = 1 + r - 2t$. The kinematic ranges for t and s are

$$\begin{aligned}\sqrt{r} &\leq t \leq \frac{1}{2}(1+r), \\ 0 &\leq s \leq (1-\sqrt{r})^2.\end{aligned}\tag{13}$$

With the T odd transverse Λ polarizations defined by

$$P_T = \frac{d\Gamma(\hat{\xi} \cdot \hat{e}_T = 1) - d\Gamma(\hat{\xi} \cdot \hat{e}_T = -1)}{d\Gamma(\hat{\xi} \cdot \hat{e}_T = 1) + d\Gamma(\hat{\xi} \cdot \hat{e}_T = -1)},$$

and from Eq. (9) we obtain

$$\begin{aligned}P_T &= \frac{3\pi}{4R_{\Lambda_b}(t)}\sqrt{s\phi}(1-s\rho^2) \\ &\times \left[(\text{Im}C_9^R C_{10}^{L*} - \text{Im}C_9^L C_{10}^{R*}) - \frac{2\hat{m}_b}{s}(1-t)(\text{Im}C_7^L C_{10}^{R*} - \text{Im}C_7^R C_{10}^{L*}) \right],\end{aligned}\tag{14}$$

where $\phi(s) = (1-r)^2 - 2s(1+r) + s^2$.

To understand Eq. (14), we first examine the hadronic currents with $V \pm A$ types of interactions. From Eq. (1), there are three possible sources which could lead to T -odd correlations:

- (i) $H_{i\mu}H_{i\nu}^\dagger L_{V(A)}^\mu L_{V(A)}^{\nu\dagger}$, $(i = 1, 2)$
- (ii) $H_{1\mu}^{L(R)}H_{2\nu}^{L(R)\dagger}L_V^\mu L_A^{\nu\dagger}$,
- (iii) $H_{1\mu}^{L(R)}H_{2\nu}^{R(L)\dagger}L_V^\mu L_A^{\nu\dagger}$,

where $H_{i\mu}^{L(R)}$ involve only left-handed (right-handed) currents. By summing all the lepton spin degrees of freedom, we get $\sum L_V^\mu L_V^{\nu\dagger} = \sum L_A^\mu L_A^{\nu\dagger} = (p_l^\mu p_{\bar{l}}^\nu + p_l^\nu p_{\bar{l}}^\mu - g^{\mu\nu} p_l \cdot p_{\bar{l}})$, which are symmetric with respect to μ and ν . Since $H_{i\mu}H_{i\nu}^\dagger \propto \varepsilon_{\mu\nu\alpha\beta} s_\Lambda^\alpha p_{\Lambda_b}^\beta$ is antisymmetric between μ and ν , it is clear that no T-odd terms can be constructed from (i). For the case in (ii), even though $\sum L_V^\mu L_A^{\nu\dagger} = -4i\varepsilon^{\mu\nu\alpha\beta} p_{l\alpha} p_{\bar{l}\beta}$ is antisymmetric, $H_{1\mu}^{L(R)}H_{2\nu}^{L(R)\dagger} \propto M_\Lambda \text{Tr}(\not{s}_\Lambda \gamma_\mu \not{p}_{\Lambda_b} \gamma_\nu) = 4M_\Lambda (s_{\Lambda\mu} p_{\Lambda_b\nu} + s_{\Lambda\nu} p_{\Lambda_b\mu} - g_{\mu\nu} s_\Lambda \cdot p_{\Lambda_b})$ is symmetric in μ and ν and thus, the possible terms in (ii) also vanish. However, T-odd correlations can arise from (iii) by observing that $H_{1\mu}^{L(R)}H_{2\nu}^{R(L)\dagger} \propto M_{\Lambda_b} \text{Tr}(\not{p}_\Lambda \not{s}_\Lambda \gamma_\mu \gamma_\nu) = 4M_{\Lambda_b} (-p_{\Lambda\mu} s_{\Lambda\nu} + p_{\Lambda\nu} s_{\Lambda\mu})$ is antisymmetric. Hence, in $\Lambda_b \rightarrow \Lambda l^+ l^-$ decays, the non-vanished T-odd terms can be induced

from $(V - A) \times (V + A)$ hadronic currents. This is, explicitly shown in Eq. (14). The TV quantity is related to $\text{Im}C_9^R C_{10}^{L*}$ and $\text{Im}C_9^L C_{10}^{R*}$, respectively. Similarly, with the dipole operators, we expect that T-odd observables are proportional to $\text{Im}C_7^R C_{10}^{L*}$ and $\text{Im}C_7^L C_{10}^{R*}$.

As seen in Eq. (14), to have a non-zero value of P_T , it is necessary to have conditions of (i) the existence of C_k^R and (ii) a phase of $C_i^L C_j^R$ ($i \neq j$). These conditions are clearly different from those of the T odd transverse lepton polarizations in both inclusive decay of $b \rightarrow sl^+l^-$ [7] and exclusive ones, such as $B \rightarrow K^{(*)}l^+l^-$ [8] and $\Lambda_b \rightarrow \Lambda l^+l^-$ [9], and other CP violating asymmetries as well [3, 10]. Such property distinguishes the T odd transverse Λ polarizations from various other T odd observables. In the standard model, since there are no C_9^R and C_{10}^R as seen from Eq. (3),

$$P_T^{SM} \propto \frac{m_s}{m_b} \text{Im} \left(C_7^{eff} C_{10} \right), \quad (15)$$

which is suppressed. We note that the contribution to P_T from the EM final state interaction is $< O(10^{-3})$. Moreover, the long-distance (LD) effects in the one-loop matrix elements of $O_{1,2}$ [4] and $\Lambda_b \rightarrow \Lambda J/\Psi$ with $J/\Psi \rightarrow l^+l^-$ are absorbed to C_9^L . On the other hand, it is clear that a large value of P_T , according to Eq. (14), can be obtained if a theory contains C_9^R or C_{10}^R or a large C_7^R , with a non-zero phase. Many theories beyond the standard model could give rise to C_9^R or C_{10}^R ; examples are the left-right symmetric and supersymmetric models.

To illustrate our result, we use SUSY models both with and without R-parity. For a SUSY model with R-parity, we take the one given by Ref. [11], denoted as $M1$, in which

$$\begin{aligned} C_7^{LM1} &= -1.75 (\delta_{23}^u)_{LL} - 0.25 (\delta_{23}^u)_{LR} - 10.3 (\delta_{23}^d)_{LR}, \\ C_9^{LM1} &= 0.82 (\delta_{23}^u)_{LR}, \\ C_{10}^{LM1} &= -9.37 (\delta_{23}^u)_{LR} + 1.4 (\delta_{23}^u)_{LR} (\delta_{33}^u)_{RL} + 2.7 (\delta_{23}^u)_{LL}, \\ C_7^{RM1} &= -10.3 (\delta_{23}^d)_{RL}, \\ C_9^{RM1} &= 1.32 (\delta_{23}^d)_{RL} (\delta_{33}^d)_{LR}, \\ C_{10}^{RM1} &= -17.6 (\delta_{23}^d)_{RL} (\delta_{33}^d)_{LR}, \end{aligned} \quad (16)$$

where the values of parameters δ_{ij}^q are as follows :

$$\begin{aligned}
(\delta_{23}^d)_{LR} &\sim (\delta_{23}^d)_{RL} \sim -3 \times 10^{-2} e^{i\theta_1}, \\
(\delta_{33}^d)_{LR} &\sim (\delta_{33}^u)_{RL} \sim 0.5, \\
(\delta_{23}^u)_{LR} &\sim -0.7 e^{i\theta_2}, \\
(\delta_{23}^u)_{LL} &\sim 0.1,
\end{aligned} \tag{17}$$

with taking the phases of θ_1 and θ_2 to be $\pi/4$ and $-\pi/10$, respectively.

To demonstrate the difference we study a SUSY model without R-parity ($M2$, see Ref. [12]). In this case we have

$$C_9^{RM2} = -C_{10}^{RM2} = \frac{\pi}{\sqrt{2}G_F\alpha_{em}} \frac{1}{V_{tb}V_{ts}^*} \frac{\lambda_{ij3}'^* \lambda_{ij2}'}{2M_{\tilde{u}_j}^2} \tag{18}$$

where

$$|\lambda_{ijk}'| \simeq \left(2\sqrt{2}G_F \tan^2 \beta\right)^{1/2} \zeta_i m_{d_j} \lambda^2 \tag{19}$$

and

$$\lambda_{ij3}'^* \lambda_{ij2}' \simeq 2\sqrt{2}G_F \tan^2 \beta \zeta_i^2 m_{d_j}^2 \lambda^4 e^{i\theta_R} \tag{20}$$

with the parameters $\lambda \sim 0.22$, $\tan \beta \sim 10$, $\zeta_i \sim 0.9$, and $\theta_R = 2\pi/5$. We note that the parameters we have chosen for the two models satisfy various experimental constraints [11, 12].

By using the parameters in Eqs. (16)-(20), the values for the form factors in Refs. [3, 6], and Eq. (10), we give the differential branching ratios (BRs) of $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ with respect to E_Λ in Figure 1 with and without including resonant states of Ψ and Ψ' , and we find that the integrated BRs for the latter are $(2.10, 1.66, 4.41) \times 10^{-6}$ for the standard and SUSY with and without R-parity models, respectively.

In Figures 2 and 3, we show $P_T(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ as a function of E_Λ/M_{Λ_b} for $M1$ and $M2$, with and without the LD contributions, respectively. As seen from the figures, even though the derivations of BRs to the standard model result are insignificant, the transverse Λ polarization asymmetries can be over 50% in both SUSY models with and without R-parity. Similar results are also expected the decay of $\Lambda_b \rightarrow \Lambda e^+ e^-$. We remark that measuring a

large P_T in $\Lambda_b \rightarrow \Lambda l^+ l^-$ is a clean indication of T violation as well as new CP violation mechanism beyond the standard model.

Finally, we note that to measure $P_T(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) \sim 10\%$ at 3σ level, at least 4.5×10^7 Λ_b decays are required if we use $BR(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) \sim 2 \times 10^{-6}$. Clearly, the measurement could be done in the second generation of B-physics experiments, such as LHCb, ATLAS, and CMS at the LHC, and BTeV at the Tevatron, which produce $\sim 10^{12} b\bar{b}$ pairs per year [13]. This is certainly within reach of a super B factory under discussion now [14].

Acknowledgments

This work was supported in part by the National Center for Theoretical Science, National Science Council of the Republic of China under Contract Nos. NSC-90-2112-M-001-069 and NSC-90-2112-M-007-040, and National Science and Engineering Research Council of Canada.

-
- [1] CPLEAR Collaboration, A. Angelopoulos et al., *Phys. Lett.* **B444**, 43 (1998).
 - [2] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
 - [3] C. H. Chen and C. Q. Geng, *Phys. Lett. B* **516**, 327 (2001) [arXiv:hep-ph/0101201].
 - [4] G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Rev. Mod. Phys.* **68**, 1230 (1996).
 - [5] T. Mannel, W. Roberts and Z. Ryzak *Nucl. Phys.* **B355**, 38 (1991);
T. Mannel and S. Recksiegel *J. Phys.* **G24**, 979 (1998).
 - [6] C.H. Chen and C. Q. Geng, *Phys Rev.* **D64**, 114024 (2001).
 - [7] F. Kruger and L.M. Sehgal, *Phys. Lett.* **B380**, 199 (1996).
 - [8] C.Q. Geng and C.P. Kao, *Phys. Rev.* **D57**, 4479 (1998).
 - [9] C. H. Chen and C. Q. Geng, *Phys. Rev. D* **64**, 074001 (2001) [arXiv:hep-ph/0106193].
 - [10] G. Hiller and A. Kagan, arXiv:hep-ph/0108074.
 - [11] F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, *Nucl. Phys.* **B477**, 321-352 (1996);
E. Lunghi, A. Masiero, I. Scimemi, and L. Silvestrini, *Nucl. Phys.* **B568**, 120-144 (2000).
 - [12] D. Guetta and E. Nardi, *Phys. Rev.* **D58**, 012001 (1998) and references therein.
 - [13] N. Harnew, *Proc. of Heavy Flavors 8*, (Southampton. 1999).
 - [14] See, *e.g.*, G. Eigen, arXiv:hep-ex/0112042.




FIG. 1: Differential BRs of $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ as a function of E_Λ with and without resonant shapes, where the solid, dash-dotted and dashed curves stand for the results of the standard and SUSY with and without R-parity models, respectively.




FIG. 2: Transverse Λ polarization in $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ as a function of E_Λ/M_{Λ_b} in the SUSY model with R-parity. The curves with and without resonant shapes represent including and no long-distance contributions, respectively.




FIG. 3: Same as Figure 2 but for the SUSY model without R-parity.